

”An effective two dimensionality” cases bring a new hope to the Kaluza-Klein[like] theories

D. Lukman and N. S. Mankoč Borštnik

Department of Physics, FMF, University of Ljubljana, Jadranska 19, 1000 Ljubljana

H. B. Nielsen

Department of Physics,

Niels Bohr Institute, Blegdamsvej 17,

Copenhagen, DK-2100

Abstract

One step towards realistic Kaluza-Klein[like] theories and a loop hole through the Witten’s ”no-go theorem” is presented for cases which we call *an effective two dimensionality* cases: In $d = 2$ the equations of motion following from the action with the linear curvature leave spin connections and zweibeins undetermined. We present the case of a spinor in $d = (1 + 5)$ compactified on a formally *infinite* disc with the *zweibein* which makes a disc curved on an almost S^2 and with the *spin connection* field which allows on such a sphere only one massless *normalizable* spinor state of a particular charge, which couples the spinor chirally to the corresponding Kaluza-Klein gauge field. We assume no external gauge fields. The masslessness of a spinor is achieved by the choice of a spin connection field (which breaks left-right symmetry), the zweibein and the normalizability condition for spinor states, which guarantee a discrete spectrum forming the complete basis. We discuss the meaning of the hole, which manifests the noncompactness of the space.

PACS numbers: 11.10.Kk, 11.25.Mj, 12.10.-g, 04.50.-h

I. INTRODUCTION

The idea of Kaluza and Klein [1] of obtaining the electromagnetism - and under the influence of their idea nowadays also the weak and colour fields [2–8, 13] - from purely gravitational degrees of freedom connected with having extra dimensions is very elegant. More than twenty five years ago the Kaluza-Klein[like] theories were studied very intensively by many authors [8, 9, 13, 14]. Although the breaking of the symmetry of the starting Lagrange density to the low energy effective ones (that is to the charges and correspondingly to the gauge fields assumed by the *standard model of the electroweak and colour interactions*) seem very promising, the idea of Kaluza and Klein was almost killed by the "no-go theorem" of E. Witten [15] telling that these kinds of Kaluza-Klein[like] theories with the gravitational fields only (that is with vielbeins and spin connections) have severe difficulties with obtaining massless fermions chirally coupled to the Kaluza-Klein-type gauge fields in $d = 1 + 3$, as required by the *standard model*. There were attempts to escape from the "no-go theorem" in compact extra spaces by having torsion [5], or by having an orbifold structure [11], or by putting extra gauge fields by hand in addition to gravity in higher dimensions [12], which is no longer the pure Kaluza-Klein[like] theory and loses accordingly the elegance.

Since there is the assumption that the space is *compact* in the "no-go theorem" of E. Witten, there are also the attempts to achieve masslessness by appropriate choices of vielbeins in *noncompact* spaces, one of works [13] is commented in the footnote [27].

There are several attempts to point out the importance of non compact extra dimensions, like [16], many of them surveyed in [17]. These attempts do not really try to keep the Kaluza-Klein approach in the original elegant version, they rather embed strings, membranes, p-branes into higher dimensional spaces. The most popular models of this kind are probably Randall-Sundrum models [18].

We are interested in this paper in extra dimensions in the Kaluza-Klein sense: that is as a possibility that the gravity (and only gravity) in extra dimensions manifests as the *standard model* gauge fields in $(1 + 3)$, coupled to the corresponding charges. In refs. [24] we achieved masslessness of spinors in the pure Kaluza-Klein[like] theory (for the case of M^{1+5} manifold broken into $M^{1+3} \times$ an infinite disc) with the appropriate choice of a boundary limiting the extra dimensions on a finite surface on a disc.

In the proposed paper we take the whole two dimensional plane, and roll it up into an

almost S^2 with one point - the south pole - excluded. It is our choice of a *zweibein* which forces the two extra dimensions into an almost S^2 . Thus, although it has a finite volume (namely the surface of S^2), the space is *non compact*. We require spinor states to be in the fifth and sixth dimensions *normalizable* [28], proving that the normalizable solutions form a complete set. It is our choice of a *particular spin connection field*, with the strengths within an interval, which allows only one normalizable massless state of a particular handedness (with respect to $(1 + 3)$), breaking the parity symmetry.

The *finite volume* of an *infinite* disc, an appropriate *choice of the spin connection field* with the strength F allowed to be within the whole interval $0 < 2F \leq 1$ and the *normalizability* requirement make the mass spectrum of our Hermitean Hamiltonian in a noncompact space discrete, with only one massless state of particular charge chirally coupled to the Kaluza-Klein gauge field. It is the sign of F which makes a choice of the handedness of a massless state, breaking the parity symmetry. The usually expected problem with extra non compact dimensions having a continuous spectrum is *not present* in our model.

For a particular choice of the strength of the spin connection field we find the states and the spectrum (the masses) analytically. This mass spectrum of states forms the complete set on our almost S^2 . For the remaining values of the strength, for all of which only one massless solution of a particular handedness in $(1 + 3)$ exists, it is not difficult to find the recursive formulas for normalizable solutions and the masses. Accordingly in this two dimensional noncompact space, with the spin connections and vielbeins which both are a part of the gravitational gauge fields and with no presence of an (additional) external field, the "no-go theorem" of E. Witten is not valid.

We also characterize the "singularity" which the spinor solutions "feel" on our infinite disc with the zweibein of a S^2 sphere, when treating the disc as the almost S^2 sphere, that is the S^2 sphere with the hole on the southern pole, so that we have almost $M^{(1+3)} \times S^2$ case, that it is almost a compact space.

Let us add: As it is not difficult to recognize, the two dimensional spaces are very special [19, 20]. Namely, in dimensions higher than two, when we have no fermions present and only the curvature in the first power in the Lagrange density, the spin connections are normally determined from the vielbein fields, and the torsion is zero. In the two dimensional spaces, the vielbeins do not determine the spin connection fields. In the present article we pay attention to cases, which we call *an effective two-dimensionality*, when the spin connections

are not fully determined by the vielbeins.

In the here proposed types of models there is the chance for having chirally mass protected fermions in a theory in which the chirally protecting effective four dimensional gauge fields are *true* Kaluza-Klein[like] fields, the degrees of which inherit from the higher dimensional gravitational ones. We are thus hoping for a revival of true Kaluza-Klein[like] models as candidates for phenomenologically viable models!

One of us has been trying for long to develop the *approach unifying spins and charges and predicting families* (N.S.M.B.) [21, 25] so that spinors which carry in $d \geq 4$ nothing but two kinds of the spin (no charges), would manifest in $d = (1 + 3)$ all the properties assumed by the *standard model* and does accordingly share with the Kaluza-Klein[like] theories the problem of masslessness of the fermions before the electroweak like types of break. We present briefly the ideas of the *approach* in the footnote [29].

Let us point out that in odd dimensional spaces and in even dimensional spaces divisible by four there is no mass protection in the Kaluza-Klein[like] theories [13, 24]. The spaces therefore, for which we can have a hope that the Kaluza-Klein[like] theories lead to chirally protected fermions and accordingly to the effective theory of the *standard model of the electroweak and colour interactions*, have $2(2n + 1)$ dimensions. And breaking symmetries in such spaces, if starting with one Weyl spinor, and accordingly with the mass protected case, should again lead to mass protected cases in accordance with the *standard model*.

II. THE ACTION, EQUATIONS OF MOTION, SOLUTIONS, PROOFS AND COMMENTS

We prove in this section that in $M^{1+3} \times$ an infinite disc with the particular zweibein and spin connection on the disc there exists only one massless normalizable (on the disc) fermion state of only one handedness and of a particular charge. It is accordingly mass protected. We also present proofs that the Hamiltonian is Hermitean and the spectra of normalizable states correspondingly discrete. For a particular strength of the spin connection field we present the spectrum and states. We discuss the properties of solutions for the strengths allowed by the normalizability requirement.

Let us first repeat the four assumptions, stressed already in the introduction.

1. We assume $2(2n + 1)$ -dimensional space, in our case $n = 1$, with only gravity, described

by the action [30]

$$\mathcal{S} = \alpha \int d^d x E \mathcal{R}. \quad (1)$$

The Riemann scalar $\mathcal{R} = \mathcal{R}_{abcd} \eta^{ac} \eta^{bd}$ is determined by the Riemann tensor $\mathcal{R}_{abcd} = f^\alpha_{[a} f^\beta_{b]} (\omega_{cd\beta, \alpha} - \omega_{ce\alpha} \omega^e_{d\beta})$, with vielbeins f^α_a [31] and the spin connections $\omega_{ab\alpha}$ (the gauge fields of $S^{ab} = \frac{i}{4}(\gamma^a \gamma^b - \gamma^b \gamma^a)$). $[a b]$ means that the antisymmetrization must be performed over the two indices a and b , E is the determinant of the inverse zweibein $e^s_\sigma, e^s_\sigma f^\sigma_t \delta^s_t$, (Eq.(2)).

2. Space M^{1+5} has the symmetry of $M^{1+3} \times$ an infinite disc with the zweibein on the disc

$$e^s_\sigma = f^{-1} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, f^\sigma_s = f \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \quad (2)$$

with

$$f = 1 + \left(\frac{\rho}{2\rho_0}\right)^2, \\ x^{(5)} = \rho \cos \phi, \quad x^{(6)} = \rho \sin \phi, \quad E = f^{-2}.$$

The last relation follows from $ds^2 = e_{s\sigma} e^s_\tau dx^\sigma dx^\tau = f^{-2}(d\rho^2 + \rho^2 d\phi^2)$. We use indices $s, t = 5, 6$ to describe the flat index in the space of an infinite plane, and $\sigma, \tau = (5), (6)$, to describe the Einstein index. ϕ determines the angle of rotations around the axis perpendicular to the disc.

3. The spin connection field is chosen to be

$$f^\sigma_{s'} \omega_{st\sigma} = iF f \varepsilon_{st} \frac{e_{s'\sigma} x^\sigma}{(\rho_0)^2}, \quad 0 < 2F \leq 1, \quad s = 5, 6, \quad \sigma = (5), (6). \quad (3)$$

4. We require normalizability of states ψ on the disc

$$\int_0^{2\pi} d\phi \int_0^\infty E \rho d\rho \psi^\dagger \psi < \infty, \quad (4)$$

as usual in quantum mechanics, allowing at most the plane waves normalized to the delta function: $\int_{-\infty}^\infty dx^{(5)} \int_{-\infty}^\infty dx^{(6)} E e^{i\vec{k}(\vec{x}-\vec{x}')} = \delta^2(\vec{x}-\vec{x}')$.

Let us make now several statements, proofs of these statements and comments, which will help to clarify the meaning of the assumptions.

Statement 1.: In the absence of the fermion fields in $d = 2$ any zweibein and any spin

connection fulfills the equations of motion.

Proof 1.: The action of Eq. (1) leads to the equations of motion [19, 21]

$$(d-2)\omega_b^c{}_c = \frac{e^a{}_\alpha}{E}\partial_\beta \left(E f^\alpha{}_{[a} f^{\beta]}{}_b \right), \quad (5)$$

which clearly demonstrate that any spin connection $\omega_b^c{}_c = \omega_b^c{}_\alpha f^\alpha{}_c$ (which can in $d=2$ have only two different indices) satisfies this equation.

Comment 1.: For $d=2$ the variation of the action (1) with respect to vielbeins leads to the equation $-e_{s\sigma}R + 4f^{\tau t}\omega_{st\sigma,\tau} = 0$, which is zero for any R ($-2R + 2R = 0$).

Statement 2.: The volume of this noncompact space (which looks almost as S^2 sphere) is finite.

Proof 2.: The volume is $\int_0^\infty f^{-2}\rho d\rho = \pi(2\rho_0)^2$.

Comments 2.: **i.)** Finite volume helps to assure the existence of normalizable spinor states on this disc. **ii.)** The symmetry of this disc, which is the symmetry of $U(1)$ group, determines the charge of spinors in $d=(1+3)$.

Statement 3.: The choice that M^{1+5} breaks into $M^{1+3} \times$ an infinite disc with no gravity in M^{1+3} and with the zweibein of Eq. (2) and the spin connection of Eq. (3) on an infinite disc makes the Lagrange density for a Weyl spinor $\mathcal{L}_W = \frac{1}{2}[(\psi^\dagger E \gamma^0 \gamma^a p_{0a} \psi) + (\psi^\dagger E \gamma^0 \gamma^a p_{0a} \psi)^\dagger]$ to be

$$\begin{aligned} \mathcal{L}_W &= \psi^\dagger \{ E \gamma^0 \gamma^n p_n + E f \gamma^0 \gamma^s \delta_s^\sigma (p_{0\sigma} + \frac{1}{2E f} \{p_\sigma, E f\}_-) \} \psi, \quad n=0,1,2,3, \\ p_{0\sigma} &= p_\sigma - \frac{1}{2} S^{st} \omega_{st\sigma}, \end{aligned} \quad (6)$$

with $E = \det(e^a{}_\alpha) = f^{-2}$, f is from Eq. (3), and with $\omega_{st\sigma}$ from Eq. (3) [32].

Proof 3.: Eq. (6) follows from the starting Lagrangean for a Weyl spinor interacting with only the vielbeins and spin connections straightforwardly.

Comment 3.: The Lagrange density of Eq. (6) assures that the Hamiltonian is Hermitean.

Statement 4.: Normalizability condition for spinors on an infinite disc curled into an almost S^2 and with the spin connection of particular choice makes a choice of a spectrum which forms a complete set.

Proof 4.: The Lagrange density of Eq. (6) leads to equations of motion (Eqs.(11,12, 15))

$$if \{ e^{i\phi 2S^{56}} \left[\frac{\partial}{\partial \rho} + \frac{i 2S^{56}}{\rho} \left(\frac{\partial}{\partial \phi} \right) - \frac{1}{2f} \frac{\partial f}{\partial \rho} (1 - 2F 2S^{56}) \right] \} \psi^{(6)} + \gamma^0 \gamma^5 m \psi^{(6)} = 0, \quad (7)$$

which look for $F=1/2$ like Legendre equations (Eq. (22)). It is the sign of F which makes a choice of the handedness of a massless state and breaks accordingly the parity symmetry.

One can prove that the only normalizable eigenstates in the interval $0 \leq \rho \leq \infty$ are those with integer parameters l and n , $(m\rho_0)^2 = l(l+1)$, in Eqs. (23). These states are Legendre polynomials and form the *complete set*. Solutions for a non integer n are singular at $\rho = 0$, while solutions with a non integer l are singular at $\rho = \infty$, both singularities make the corresponding eigenstates not normalizable.

Comments 4.: i.) In the subsection II A of this section the solutions of Eq. (7) are discussed for any choice of F in the interval $0 < 2F \leq 1$. All the normalizable solutions can for any F in this interval be expressed as a normalizable superposition of a complete set of Legendre polynomials and have the discrete spectrum. *ii.)* In the limit when $\rho_0 \rightarrow \infty$, f (in Eq. (15, next section) goes to one and the two equations, Eq. (15), define the recurrence relations between the Bessel functions of an integer order ($\mathcal{A}_n(\rho m) = J_n(\rho m)$ and $\mathcal{B}_{n+1}(\rho m) = J_{n+1}(\rho m)$) for any mass m . Making the limit $\rho_0 \rightarrow \infty$ in Eq. (22) in next section, with the discrete mass term $(m\rho_0)^2 = l(l+1)$ one again reproduces the Bessel equation, if putting $l = m\rho_0$. (Bessels functions can be squared normalized only within a finite radius, determined by zeros.) With ρ_0 going to infinity the distance between m -values solving this constraint goes to zero, so that in this limit the system of allowed m values approaches the continuum (all m values). This is satisfactory because this limit corresponds to our already non-compact space approaches, the usual flat two-dimensional space (with which one would have a truly fully 5 +1 dimensional world in which of course the spectrum seen as (3 + 1)-dimensional one should be continuous). *iii.)* For any finite ρ_0 can the plane wave in the fifth and sixth dimension be expressed in terms of the Legendre polynomials. To a plane wave in general many Legendre polynomials contribute, each corresponding to a different mass. There is the solution for $2F = 1$ which is independent of x^σ , $\sigma \in \{(5), (6)\}$. It corresponds to massless solution. This solution can be called the plane wave with momentum zero. In the limit $\rho_0 \rightarrow \infty$ the definition for the plane waves in flat space follows. *Statement 5.: The zweibein (Eq.(2)) and the spin connection (Eq.(3)) with the parameter F within the interval $0 < 2F \leq 1$ allow only one massless spinor of a particular charge.*

Proof 5.: It is proven in the next subsection, in the last paragraph before Eq. (11), that it is the term $\psi^\dagger E f \gamma^0 \gamma^s \delta_s^\sigma (p_{0\sigma} + \frac{1}{2Ef} \{p_\sigma, E f\}_-) \psi$ in the Lagrange density (Eq.(6)), which manifests as the mass term m in Eq. (7). There is a term in Eq. (7), namely $-i f e^{i\phi 2S^{56}} \frac{1}{2f} \frac{\partial f}{\partial \rho} (1 - 2F 2S^{56}) \psi^{(6)}$, which clearly distinguishes between the two possible values of the spin operator S^{56} in $d = 5, 6$, when this term applies on the state $\psi^{(6)}$, dis-

tinguishing correspondingly also between the two possible handedness of the state $\psi^{(6)}$ in $d = (1 + 3)$. It is shown in the next subsection that a normalizable massless state ($m = 0$ in Eq. (7)) must fulfil the condition: $(0 \leq (1 - 2F 2S^{56}) < 1) \psi^{(6)}$. The sign of F chooses the handedness of a massless normalizable spinor state.

Comments 5. i.) Having the rotational symmetry around the axis perpendicular to the plane of the fifth and the sixth dimension it is meaningful to require that $\psi^{(6)}$ is the eigenfunction of the total angular momentum operator ($M^{56} = x^5 p^6 - x^6 p^5 + S^{56}$) in the fifth and sixth dimension $M^{56} = (-i \frac{\partial}{\partial \phi} + S^{56})$; $M^{56} \psi^{(6)} = (n + \frac{1}{2}) \psi^{(6)}$ (Eqs.(13,14, 12)). *ii.)* The only massless state, which fulfills the normalization condition (see Eq.(18)) for a positive F , is the state with the property $2S^{56} \psi^{(6)} = \psi^{(6)}$. Its charge (spin on the disc) is for $0 < 2F \leq 1$ equal to $\frac{1}{2}$ as it is shown in section IV. *iii.)* All the other states are massive. *iv.)* The current in the radial direction is for all these cases equal to zero for any F .

Detailed derivations of equations of motion and solutions are presented in subsection II A of this section.

Let us summarize this section. We have a Weyl spinor in $d = (1 + 5)$ -dimensional space. This space breaks into M^{1+3} cross an infinite disc with the zweibein which formally looks almost – up to a hole in the southern pole – as a S^2 sphere, while a chosen spin connection allows on such an infinite disc only one normalizable massless state. The *Hamiltonian is Hermitean*, the mass spectrum of *normalizable* states is correspondingly discrete and the probability for a fermion to escape out of the disc is zero [33].

Allowing the whole interval of the strength of the spin connection fields ($0 < 2F \leq 1$) the spin connection field is not fine tuned. For a particular choice of the constant of the spin connection field, that is for $2F = 1$, the normalizable solutions are expressible with the Legendre polynomials and the massive states manifest a spectrum $m\rho_0 = l(l + 1)$, with $l = 0, 1, 2, \dots$ and $-l \leq n \leq 1$. $n + 1/2$ is the charge of the spectrum.

A free choice of a zweibein and a spin connection field in the action of Eq. (1) is possible only in $d = 2$ dimensional spaces (the presence of fermions might make this possible also for $d > 2$).

Let us point out that the "two dimensionality" can be simulated in any dimension larger than two, if vielbeins and spin connections are completely flat in all but two dimensions (this point is discussed also in the ref. [13]).

A. Solutions of the equations of motion for spinors

We look for the solutions of the equations of motion (6) for a spinor in $(1+5)$ -dimensional space, which breaks into $M^{(1+3)} \times$ an infinite disc curved into a noncompact "almost" S^2 sphere as a superposition of all four $(2^{6/2-1})$ states of a single Weyl representation. (We kindly ask the reader to see the technical details about how to write a Weyl representation in terms of the Clifford algebra objects after making a choice of the Cartan subalgebra, for which we take: S^{03}, S^{12}, S^{56} in the refs. [25].) In our technique one spinor representation—the four states, which all are the eigenstates of the chosen Cartan subalgebra with the eigenvalues $\frac{k}{2}$, correspondingly—are the following four products of projectors $\overset{ab}{[k]}$ and nilpotents $\overset{ab}{(k)}$:

$$\begin{aligned}\varphi_1^1 &= \overset{56}{(+)}\overset{03}{(+i)}\overset{12}{(+)} \psi_0, \\ \varphi_2^1 &= \overset{56}{(+)}\overset{03}{[-i]}\overset{12}{[-]} \psi_0, \\ \varphi_1^2 &= \overset{56}{[-]}\overset{03}{[-i]}\overset{12}{(+)} \psi_0, \\ \varphi_2^2 &= \overset{56}{[-]}\overset{03}{(+i)}\overset{12}{[-]} \psi_0,\end{aligned}\tag{8}$$

where ψ_0 is a vacuum state for the spinor state. If we write the operators of handedness in $d = (1+5)$ as $\Gamma^{(1+5)} = \gamma^0 \gamma^1 \gamma^2 \gamma^3 \gamma^5 \gamma^6 (= 2^3 i S^{03} S^{12} S^{56})$, in $d = (1+3)$ as $\Gamma^{(1+3)} = -i \gamma^0 \gamma^1 \gamma^2 \gamma^3 (= 2^2 i S^{03} S^{12})$ and in the two dimensional space as $\Gamma^{(2)} = i \gamma^5 \gamma^6 (= 2 S^{56})$, we find that all four states are left handed with respect to $\Gamma^{(1+5)}$, with the eigenvalue -1 , the first two states are right handed and the second two states are left handed with respect to $\Gamma^{(2)}$, with the eigenvalues 1 and -1 , respectively, while the first two are left handed and the second two right handed with respect to $\Gamma^{(1+3)}$ with the eigenvalues -1 and 1 , respectively. Taking into account Eq. (8) we may write the most general wave function $\psi^{(6)}$ obeying Eq. (7) in $d = (1+5)$ as

$$\psi^{(6)} = \mathcal{A} \overset{56}{(+)} \psi_{(+)}^{(4)} + \mathcal{B} \overset{56}{[-]} \psi_{(-)}^{(4)},\tag{9}$$

where $\mathcal{A}$ and $\mathcal{B}$ depend on x^σ , while $\psi_{(+)}^{(4)}$ and $\psi_{(-)}^{(4)}$ determine the spin and the coordinate dependent parts of the wave function $\psi^{(6)}$ in $d = (1+3)$

$$\begin{aligned}\psi_{(+)}^{(4)} &= \alpha_+ \overset{03}{(+i)} \overset{12}{(+)} + \beta_+ \overset{03}{[-i]} \overset{12}{[-]}, \\ \psi_{(-)}^{(4)} &= \alpha_- \overset{03}{[-i]} \overset{12}{(+)} + \beta_- \overset{03}{(+i)} \overset{12}{[-]}.\end{aligned}\tag{10}$$

Using $\psi^{(6)}$ in Eq. (7) and separating dynamics in $(1+3)$ and on the infinite disc the following relations follow, from which we recognize the mass term m : $\frac{\alpha_+}{\alpha_-} (p^0 - p^3) - \frac{\beta_+}{\beta_-} (p^1 - ip^2) = m$,

$\frac{\beta_+}{\beta_-}(p^0+p^3)-\frac{\alpha_+}{\beta_-}(p^1+ip^2)=m$, $\frac{\alpha_-}{\alpha_+}(p^0+p^3)+\frac{\beta_-}{\alpha_+}(p^1-ip^2)=m$, $\frac{\beta_-}{\beta_+}(p^0-p^3)+\frac{\alpha_-}{\beta_+}(p^1-ip^2)=m$. One notices that for massless solutions ($m=0$) $\psi_{(+)}^{(4)}$ and $\psi_{(-)}^{(4)}$ decouple. Taking the above derivation into account Eq. (7) transforms into

$$f \{ (p_{05} + i2S^{56} p_{06}) + \frac{1}{2E} \{ p_5 + i2S^{56} p_6, Ef \}_- \} \psi^{(6)} + \gamma^0 \gamma^5 m \psi^{(6)} = 0. \quad (11)$$

For $x^{(5)}$ and $x^{(6)}$ from Eq. (3) and for the zweibein from Eqs.(2,3) and the spin connection from Eq.(3) one obtains

$$if \{ e^{i\phi 2S^{56}} [\frac{\partial}{\partial \rho} + \frac{i2S^{56}}{\rho} (\frac{\partial}{\partial \phi}) - \frac{1}{2f} \frac{\partial f}{\partial \rho} (1 - 2F 2S^{56})] \} \psi^{(6)} + \gamma^0 \gamma^5 m \psi^{(6)} = 0. \quad (12)$$

Having the rotational symmetry around the axis perpendicular to the plane of the fifth and the sixth dimension we require that $\psi^{(6)}$ is the eigen function of the total angular momentum operator $M^{56} = x^5 p^6 - x^6 p^5 + S^{56} = -i \frac{\partial}{\partial \phi} + S^{56}$

$$M^{56} \psi^{(6)} = (n + \frac{1}{2}) \psi^{(6)}. \quad (13)$$

Accordingly we write

$$\psi^{(6)} = \mathcal{N} (\mathcal{A}_n \overset{56}{(+)} \psi_{(+)}^{(4)} + \mathcal{B}_{n+1} e^{i\phi} \overset{56}{[-]} \psi_{(-)}^{(4)}) e^{in\phi}. \quad (14)$$

After taking into account that $S^{56} \overset{56}{(+)} = \frac{1}{2} \overset{56}{(+)}$, while $S^{56} \overset{56}{[-]} = -\frac{1}{2} \overset{56}{[-]}$ we end up with the equations of motion for $\mathcal{A}_n$ and $\mathcal{B}_{n+1}$ as follows

$$\begin{aligned} -if \{ (\frac{\partial}{\partial \rho} + \frac{n+1}{\rho}) - \frac{1}{2f} \frac{\partial f}{\partial \rho} (1 + 2F) \} \mathcal{B}_{n+1} + m \mathcal{A}_n &= 0, \\ -if \{ (\frac{\partial}{\partial \rho} - \frac{n}{\rho}) - \frac{1}{2f} \frac{\partial f}{\partial \rho} (1 - 2F) \} \mathcal{A}_n + m \mathcal{B}_{n+1} &= 0. \end{aligned} \quad (15)$$

Let us treat first the massless case ($m=0$). Taking into account that $F \frac{f-1}{f\rho} = \frac{\partial}{\partial \rho} \ln f^{\frac{F}{2}}$ and that $E = f^{-2}$, it follows

$$\begin{aligned} \frac{\partial \ln(\mathcal{B}_n \rho^n f^{-F-1/2})}{\partial \rho} &= 0, \\ \frac{\partial \ln(\mathcal{A}_n \rho^{-n} f^{F-1/2})}{\partial \rho} &= 0. \end{aligned} \quad (16)$$

We get correspondingly the solutions

$$\begin{aligned} \mathcal{B}_n e^{in\phi} &= \mathcal{B}_0 e^{in\phi} \rho^{-n} f^{F+1/2}, \\ \mathcal{A}_n e^{in\phi} &= \mathcal{A}_0 e^{in\phi} \rho^n f^{-F+1/2}. \end{aligned} \quad (17)$$

Requiring that only normalizable (square integrable) solutions are acceptable

$$\begin{aligned} 2\pi \int_0^\infty E \rho d\rho \mathcal{A}_n^* \mathcal{A}_n &< \infty, \\ 2\pi \int_0^\infty E \rho d\rho \mathcal{B}_n^* \mathcal{B}_n &< \infty, \end{aligned} \quad (18)$$

it follows

$$\begin{aligned} \text{for } \mathcal{A}_n : -1 < n < 2F, \\ \text{for } \mathcal{B}_n : 2F < n < 1, \quad n \text{ is an integer.} \end{aligned} \quad (19)$$

One immediately sees that for $F = 0$ there is no solution for the zweibein from Eq. (3).

Eq. (19) tells us that the strength F of the spin connection field $\omega_{56\sigma}$ can make a choice between the two massless solutions $\mathcal{A}_n$ and $\mathcal{B}_n$: For

$$0 < 2F \leq 1 \quad (20)$$

the only massless solution is the left handed spinor with respect to $(1 + 3)$

$$\psi_{\frac{1}{2}}^{(6)m=0} = \mathcal{N}_0 f^{-F+1/2} \binom{56}{+} \psi_{(+)}^{(4)}. \quad (21)$$

It is the eigen function of M^{56} with the eigenvalue $1/2$. No right handed massless solution is allowed. For the particular choice $2F = 1$ the spin connection field $-S^{56}\omega_{56\sigma}$ compensates the term $\frac{1}{2Ef}\{p_\sigma, Ef\}_-$ and the left handed spinor with respect to $d = (1 + 3)$ becomes a constant with respect to ρ and ϕ .

For $2F = 1$ it is easy to find also all the massive solutions of Eq. (15). Introducing $u = \frac{\rho}{2\rho_0}$ and assuming that $2F = 1$ one finds from Eq. (15)

$$\begin{aligned} \mathcal{B}_{n+1} &= \frac{i}{2\rho_0 m} (1 + u^2) \left(\frac{d}{du} - \frac{n}{u} \right) \mathcal{A}_n^m, \\ \left\{ \left(\frac{1 + u^2}{2} \right)^2 \left(\frac{d^2}{du^2} + \frac{1}{u} \frac{d}{du} - \frac{n^2}{u^2} \right) + (\rho_0 m)^2 \right\} \mathcal{A}_n^m &= 0. \end{aligned} \quad (22)$$

If one expresses $(\frac{\rho}{2\rho_0})^2 = \frac{1-x}{1+x}$, with $-1 \leq x \leq 1$ for $0 \leq \rho \leq \infty$, it follows that $f = \frac{2}{1+x}$, $\frac{dx}{du} = \frac{-4u}{(1+u^2)^2}$ and $\frac{4u^2}{(1+u^2)^2} = (1-x^2)$. Then Eq. (22) transforms into the equations of motion for the associate Legendre polynomials $\mathcal{A}_n^{(\rho_0 m)^2=l(l+1)} = P_n^l$, if we assume that $(\rho_0 m)^2 = l(l+1)$

$$\begin{aligned} \left((1-x^2) \frac{d^2}{dx^2} - 2x \frac{d}{dx} - \frac{n^2}{1-x^2} + l(l+1) \right) \mathcal{A}_n^{(\rho_0 m)^2=l(l+1)} &= 0, \\ l(l+1) &= (\rho_0 m)^2, \\ \mathcal{B}_{n+1}^{(\rho_0 m)^2=l(l+1)} &= \frac{-i}{\rho_0 m} \sqrt{1-x^2} \left(\frac{d}{dx} + \frac{n}{1-x^2} \right) \mathcal{A}_n^{(\rho_0 m)^2=l(l+1)}. \end{aligned} \quad (23)$$

From the above equations we see that for $m = 0$, that is for the massless case, the only solution with $n = 0$ exists, which is $\mathcal{A}_0^{(\rho_0 m)^2=0}$, which is a constant (in agreement with our discussions above).

It is not difficult to prove that there is no normalizable solutions of Eq. (23) for an arbitrary $m \rho_0$, which is not of the kind $(m \rho_0)^2 = l(l+1)$, with l an integer and also not for a noninteger n . The solutions of Eq. (23) are, namely, not square integrable on the interval $-1 \leq x \leq 1$ for $l \neq$ an integer, and $\nu \neq$ an integer. $P_\nu^n(x \rightarrow -1+0)$ are unbounded, going to ∞ , while they are bounded at $(x \rightarrow 1-0)$. One also finds that $P_n^\mu \rightarrow \infty$, if $(x \rightarrow 1-0)$, unless $\mu = \pm m$, with m an integer. (See ref. [26], sect. 5.18, pages 255-258.)

Accordingly the massive solutions with the masses equal to $m = l(l+1)/\rho_0$ (we use the units in which $c = 1 = \hbar$) and the eigenvalues of M^{56} ((Eq. 13))—which is the charge as we see in section IV—equal to $(\frac{1}{2} + n)$, with $-l \leq n \leq l$, $l = 1, 2, \dots$, are

$$\psi_{n+1/2}^{(6)(\rho_0 m)^2=l(l+1)} = \mathcal{N}_{n+1/2}^l \left(\binom{56}{+} \psi_{(+)}^{(4)} + \frac{i}{2\sqrt{l(l+1)}} \binom{56}{-} \psi_{(-)}^{(4)} e^{i\phi} (1+u^2) \left(\frac{d}{du} - \frac{n}{u} \right) \right) \cdot e^{in\phi} \mathcal{A}_n^{(\rho_0 m)^2=l(l+1)}, \quad (24)$$

with $\mathcal{A}_n^{(\rho_0 m)^2=l(l+1)}(x)$, which are the associate Legendre polynomials $P_n^l(x)$, where $x = \frac{1-u^2}{1+u^2}$, and $u = \frac{\rho}{2\rho_0}$ [34]. It is not difficult to see that the solutions of Eq. (15) for $2F = 1$, $\psi_{\frac{1}{2}}^{(6)m=0}$ and $\psi_{n+1/2}^{(6)(\rho_0 m)^2=l(l+1)}$, are normalizable on the infinite disc curved into almost S^2 ($2\pi \int \rho d\rho E \psi_{n+1/2}^{(6)(\rho_0 m)^2=l(l+1)\dagger} \psi_{n+1/2}^{(6)(\rho_0 m)^2=l(l+1)} < \infty$, with $E = f^{-2}$). One can show as well that the eigenstates, with the discrete eigenvalues $(\rho_0 m)^2 = l(l+1)$, are orthogonal ($\int d^2x E (\psi_{n'+1/2}^{(6)(\rho_0 m)^2=l'(l'+1)\dagger} \psi_{n+1/2}^{(6)(\rho_0 m)^2=l(l+1)}) = \delta^{ll'} \delta^{nn'} \propto \int d^2x e^{-i(n'-n)\phi} \{\mathcal{B}_{n'+1}^{l'+} \mathcal{B}_{n+1}^l + \mathcal{A}_{n'}^{l'+} \mathcal{A}_n^l\}$) for all pairs of $(l, n), (l', n')$, the spectrum is obviously discrete as it should be for the Hermitean Hamiltonian with the boundary conditions determined by normalizability of states.

To find solutions for all F in the interval $0 < F \leq \frac{1}{2}$, besides the massless one $\psi_{\frac{1}{2}}^{(6)m=0}$, is a more tough work. Yet one can expect that on the space of normalizable functions the Hamiltonian will stay Hermitean and since an infinitesimal change of the constant F from $F = \frac{1}{2}$ to a tiny smaller F can not spoil the discreteness of the Hamiltonian eigenvalues, the spectrum would stay discrete. One can see that the current in the radial direction is zero for any F . We studied these solutions and found the discrete spectrum, a paper is in

preparation.

(Let us recognize that $e^{in\phi} P_n^l$ are spherical harmonics Y_n^l . Expressing ρ with ϑ , $\frac{\rho}{2\rho_0} = \sqrt{\frac{1-\cos\vartheta}{1+\cos\vartheta}}$ we rewrite the equations of motion (Eq.15) as follows

$$\begin{aligned} & \left(\frac{\partial}{\partial\vartheta} + \frac{n+1 - (F+1/2)(1-\cos\vartheta)}{\sin\vartheta} \right) \mathcal{B}_{n+1} + i\rho_0 m \mathcal{A}_n = 0, \\ & \left(\frac{\partial}{\partial\vartheta} + \frac{-n + (F-1/2)(1-\cos\vartheta)}{\sin\vartheta} \right) \mathcal{A}_n + i\rho_0 m \mathcal{B}_{n+1} = 0. \end{aligned} \quad (25)$$

III. SINGULARITIES ON AN ALMOST S^2 SPHERE

In this section we comment on singularities "felt" by a spinor if a noncompact disc with the zweibein from Eq. (2) and the spin connections from Eq. (3) is understood as the S^2 sphere with a hole on the southern pole.

Intuitively it is not difficult to see that we are in troubles if we want the chiral fermion field of Eq. (21), that is $\psi_{\frac{1}{2}}^{(6)m=0} = \mathcal{N}_0 f^{-F+1/2} \binom{56}{+} \psi_{(+)}^{(4)}$, on a two dimensional space to be an eigenstate of some rotational operator M^{56} , if the two dimensional space has to have the topology of S^2 , while the spin of the fermion contributes to M^{56} in the "usual way"

$$M^{56} = S^{56} + K^{56}, \quad (26)$$

where K^{56} is the Killing vector, like in Eq. (13) ($K^{56} = x^5 p^6 - x^6 p^5$). Near the starting point (the origin, the northern pole of S^2) on the topologically S^2 sphere the Killing operator functions as the orbital angular momentum ($L^{56} = x^{(5)} p^{(6)} - x^{(6)} p^{(5)}$) and has to be added to the spin part S^{56} , just as it is in the flat two-dimensional space. Going away from the starting point the action of M^{56} may be more complicated as just a simple sum in Eq. (26). Because of the S^2 topology there has to be namely yet another point at which the orbital Killing generator eigenvalue goes to zero, since there has to be a point, the south pole, which is left invariant under the orbital Killing transportation as it is at the starting point, at the north pole.

It is also easy to see that on the two-dimensional S^2 , the orientation of the Killing transportation in the infinitesimal neighbourhood of this second stable point, the south pole, is in the *opposite* direction with respect to the orientation of the Killing transportation around the north pole.

If we want to have on S^2 only a spinor of one handedness, let say the spinor $\psi_{\frac{1}{2}}^{(6)m=0}$ of Eq. (21), then we should count at the south pole the orbital symmetry generator with


the opposite sign relative to S^{56} as we do at the starting point (see Eqs. (45,43)). In order to be able to have on the two-dimensional S^2 surface a spinor of only one handedness, we have to let the phase rotation generated by S^{56} part of M^{56} relative to the Killing part at the south pole to be of the opposite sign with respect to the north pole. Namely, when we consider smaller and smaller circles around the south pole, the phase of the single handedness spinor state must be rotated under M^{56} so that when extrapolating to the south pole the phase rotation correspond to the spin, which is inverted relative to the orientation of the two-dimensional space of the S^2 surface.

Therefore, embedding the S^2 sphere into a three-dimensional Euclidean space, it is not surprising that if we want a spinor of one handedness and succeed to implement it at the north pole in an outward normal direction, we can hardly implement it at the south pole. We might hope for the compensation by the orbital part of M^{56} , except at the poles. This means that we could have a state of a handed spinor if the wave function goes to zero at at least one of the poles, say the southern pole (see Eqs. (21,19)).

A. Formal introduction of a singular point

We might formally introduce at the south pole a special singularity, so that we require the wave function instead to behave at the south pole in the usual differentiable way, to be differentiable only after being multiplied (corrected) by a phase factor: Instead of ψ we require that $e^{i\phi^{SP}} \psi$ is our wave differentiable function in the neighbourhood of the singular point at the south pole, the phase factor $e^{i\phi^{SP}}$ itself behaving singularly. By making this modified requirement of the differentiability we effectively change the orbital angular momentum of the wave function by one unit of $\hbar$ before we require the wave function to be smooth or differentiable. Thereby we have made the requirement that the actual wave function should have a rather unphysical extra bit of a negative angular momentum around the south pole. We must admit that it looks rather strange from the physical point of view, unless we recognize that this smoothness condition is to simulate the non-compactness of the S^2 space, which only after adding a singular point becomes an S^2 at all.

When changing the differentiability of the wave function in the neighbourhood of the singular point with the requirement that the wave function must be multiplied by a phase, we recognize that such a phase multiplication of the wave function appears when transforming

FIG. 1: Transforming coordinates from the north to the south pole on S^2 .

the coordinate system from the northern to the southern pole, as we can see in equation (39) below. This phase transformation of the wave function requires the appearance of the spin connection field, as can be seen in Eq.(35): The gauge transformation of any spin connection field (when transforming the coordinate system), appears even if the spin connection field is zero and manifests in the second term of this equation.

B. Gauge transformations from the northern to the southern pole

To demonstrate further what does the hole do in the noncompact space of an almost S^2 sphere let us transform the coordinate system from the northern to the southern pole of the sphere S^2 as the S^2 would be a sphere made out of an infinite plane with the zweibein of a sphere and look at how do the equations of motion and the wave functions transform correspondingly and how do they demonstrate the noncompactness of our space.

From Fig. 1 we read

$$x^{NP(5)} = \left(\frac{2\rho_0}{\rho^{SP}}\right)^2 x^{SP(5)}, \quad x^{NP(6)} = -\left(\frac{2\rho_0}{\rho^{SP}}\right)^2 x^{SP(6)}, \quad (27)$$

and

$$\rho^{SP} \rho^{NP} = (2\rho_0)^2, \quad E^{NP} d^2 x^{NP} = E^{SP} d^2 x^{SP}, \quad (28)$$

where $x^{NP\sigma}, \sigma = (5), (6)$ stay for up to now used $x^\sigma, \sigma = (5), (6)$, while $x^{SP\sigma}, \sigma = (5), (6)$ stay for coordinates when we put our coordinate system at the southern pole and ρ_0 is the radius

of S^2 as before. We have $E^{SP} = (1 + (\frac{\rho^{SP}}{2\rho_0})^2)^{-2}$ and $E^{NP} = (1 + (\frac{\rho^{NP}}{2\rho_0})^2)^{-2} = (\frac{2\rho_0}{\rho^{SP}})^4 E^{SP}$. We also can write $x^{NP\sigma} = (\frac{2\rho_0}{\rho^{SP}})^2 (-)^{1+\sigma} x^{SP\sigma}$.

We ought to transform the Lagrange density (Eq.(6)) expressed with respect to the coordinates at the northern pole

$$\begin{aligned}\mathcal{L}_W^{NP} &= \psi^{NP\dagger} E^{NP} \gamma^0 \gamma^s (f^{NP\sigma}{}_s p_{0\sigma}^{NP} + \frac{1}{2E^{NP}} \{p_\sigma^{NP}, E^{NP} f^{NP\sigma}{}_s\}_-) \psi^{NP}, \\ p_{0\sigma}^{NP} &= p_\sigma^{NP} - \frac{1}{2} S^{st} \omega_{st\sigma}^{NP}, \\ f^{NP\sigma}{}_s \omega_{s't'\sigma}^{NP} &= \frac{iF \delta_s^\sigma \varepsilon_{s't'} x_\sigma^{NP}}{\rho_0^2}\end{aligned}\tag{29}$$

to the corresponding Lagrange density $\mathcal{L}_W^{SP}$ expressed with respect to the coordinates at the southern pole by assuming

$$\psi^{NP} = S \psi^{SP}.\tag{30}$$

We use the antisymmetric tensor $\varepsilon^{(5)(6)} = 1 = -\varepsilon^{(5)}_{(6)}$. We recognize that

$$\begin{aligned}f^{NP\sigma}{}_s &= f^{SP\sigma'}{}_t \frac{\partial x^{NP\sigma}}{\partial x^{SP\sigma'}} O^{-1t}{}_s, \\ f^{SP\sigma}{}_s &= f^{SP} \delta_s^\sigma, \quad f^{SP} = (1 + (\frac{\rho^{SP}}{2\rho_0})^2).\end{aligned}\tag{31}$$

The matrix O takes care that the zweibein expressed with respect to the coordinate system at the southern pole is diagonal: $f^{SP\sigma}{}_s = f^{SP} \delta_s^\sigma$

$$O = \begin{pmatrix} -\cos(2\phi + \pi) & -\sin(2\phi + \pi) \\ \sin(2\phi + \pi) & -\cos(2\phi + \pi) \end{pmatrix}.\tag{32}$$

Requiring that

$$S^{-1} \gamma^0 \gamma^s S O^{-1t}{}_s = \gamma^0 \gamma^t,\tag{33}$$

from where it follows that $S^{-1} S^{st} S O^{-1s'}{}_s O^{-1t'}{}_t = S^{s't'}$, and recognizing that $p_\sigma^{NP} = \frac{\partial x^{SP\sigma'}}{\partial x^{NP\sigma}} p_{\sigma'}^{SP}$, with $p_\sigma^{SP} = i \frac{\partial}{\partial x^{SP\sigma}}$, we find that $\gamma^s f^{NP\sigma}{}_s p_{0\sigma}^{NP} (= \gamma^s f^{NP\sigma}{}_s (p_\sigma^{NP} - \frac{1}{2} S^{st} \omega_{st\sigma}^{NP}))$ transforms into $\gamma^s f^{SP\sigma}{}_s p_{0\sigma}^{SP}$

$$\begin{aligned}\gamma^s f^{SP\sigma}{}_s p_{0\sigma}^{SP} &= \gamma^s f^{SP\sigma}{}_s \{p_\sigma^{SP} - \\ &\quad \frac{1}{2} S^{s't'} i \varepsilon_{s't'} (\frac{F x_\sigma^{SP}}{f^{SP} (f^{SP} - 1) \rho_0^2} + 2i \frac{\varepsilon_\sigma{}^\tau x_\tau^{SP}}{(2\rho_0)^2 (f^{SP} - 1)})\}.\end{aligned}\tag{34}$$

In the above equation we took into account that $\omega_{st\sigma}^{NP}$ transforms into $O^{-1s'}_s O^{-1t'}_t \frac{\partial x^{SP\sigma'}}{\partial x^{NP\sigma}} (\omega_{s't'\sigma'}^{NP} + O_{s't''}(\frac{\partial}{\partial x^{NP\sigma''}} O^{-1t''}_{t'}) \frac{\partial x^{NP\sigma''}}{\partial x^{SP\sigma'}})$, from where it follows that $\omega_{st\sigma}^{NP}$ transforms into

$$\begin{aligned} & O^{-1s'}_s O^{-1t'}_t \frac{\partial x^{SP\sigma'}}{\partial x^{NP\sigma}} \omega_{s't'\sigma'}^{SP}, \\ \omega_{st\sigma}^{SP} = & i\varepsilon_{st} \left\{ \frac{F x_\sigma^{SP}}{f^{SP} \rho_0^2 (f^{SP} - 1)} + 2i \frac{\varepsilon_\sigma^\tau x_\tau^{SP}}{(2\rho_0)^2 (f^{SP} - 1)} \right\}. \end{aligned} \quad (35)$$

Similarly we transform the term $\gamma^s \frac{1}{2E^{NP}} \{p_\sigma^{NP}, E^{NP} f^{NP\sigma}_s\}_-$ into

$$\gamma^s \left(\frac{1}{2E^{SP}} \{p_\sigma^{SP}, E^{SP} f^{SP\sigma}_s\}_- + \frac{1}{2} f^{SP\sigma}_s \{p_\sigma^{SP}, \ln(\frac{\rho^{SP}}{2\rho_0})^2\}_- \right). \quad (36)$$

The action $\int d^2x^{NP} \mathcal{L}_W^{NP}$, with the density from Eq.(6), transforms, when the coordinate system is put at the southern pole, as follows

$$\begin{aligned} \int d^2x^{NP} \mathcal{L}_W^{NP} = & \int d^2x^{SP} \psi^{SP\dagger} E^{SP} S^\dagger \gamma^0 \gamma^s (f^{SP\sigma'}_t \frac{\partial x^{NP\sigma}}{\partial x^{SP\sigma'}} O^{-1t}_s \frac{\partial x^{SP\sigma''}}{\partial x^{NP\sigma}} p_{0\sigma''}^{SP} + \\ & \frac{1}{2E^{SP}} \{p_\sigma^{SP}, E^{SP} f^{SP\sigma}_s\}_- + \frac{1}{2} f^{SP\sigma}_s \{p_\sigma^{SP}, \ln(f^{SP} - 1)\}_-) S \psi^{SP}, \end{aligned} \quad (37)$$

which leads to the Lagrange density

$$\begin{aligned} \mathcal{L}_W^{SP} = & \psi^{SP\dagger} E^{SP} \gamma^0 \gamma^s (f^{SP\sigma}_s p_{0\sigma}^{SP} + \\ & \frac{1}{2E^{SP}} \{p_\sigma^{SP}, E^{SP} f^{SP\sigma}_s\}_- + \frac{1}{2} f^{SP\sigma}_s \{p_\sigma^{SP}, \ln(\frac{\rho^{SP}}{2\rho_0})^2\}_-) \psi^{SP}. \end{aligned} \quad (38)$$

The requirement that $S^{-1} \gamma^0 \gamma^s S O^{-1t}_s = \gamma^0 \gamma^t$ is fulfilled by the operator $S = e^{-iS^{56}\omega_{56}}$, and $\omega_{56} = 2\phi + \pi$, so that in the space of the two vectors $(\begin{smallmatrix} 56 \\ (+) \end{smallmatrix} \psi_{(+)}^{(4)}, \begin{smallmatrix} 56 \\ [-] \end{smallmatrix} \psi_{(-)}^{(4)})$

$$S = \begin{pmatrix} e^{i(\phi^{NP} + \frac{\pi}{2})} & 0 \\ 0 & e^{-i(\phi^{NP} + \frac{\pi}{2})} \end{pmatrix}, \quad (39)$$

with $\phi^{NP} = -\phi^{SP}$, while we have

$$\gamma^0 \gamma^5 = \begin{pmatrix} 0 & -1 \\ -1 & 0 \end{pmatrix}, \gamma^0 \gamma^6 = \begin{pmatrix} 0 & i \\ -i & 0 \end{pmatrix}. \quad (40)$$

Let us look how does an eigenstate of M^{ab} from Eq. (13), expressed with respect to the coordinate at the northern pole

$$\psi_{n+\frac{1}{2}}^{NP(6)} = (\alpha_n(\rho^{NP}) \begin{smallmatrix} 56 \\ (+) \end{smallmatrix} \psi_{(+)}^{(4)} + i\beta_n(\rho^{NP}) \begin{smallmatrix} 56 \\ [-] \end{smallmatrix} \psi_{(-)}^{(4)} e^{i\phi^{NP}}) e^{in\phi^{NP}}, \quad (41)$$

with the property

$$M^{NP56} \psi_{n+\frac{1}{2}}^{NP(6)} = (n + \frac{1}{2}) \psi_{n+\frac{1}{2}}^{NP(6)}, \quad (42)$$

where $M^{NP56} = (S^{56} - i \frac{\partial}{\partial \phi^{NP}})$, look like when we put the coordinate system at the southern pole. When putting the coordinate system at the southern pole not only ϕ^{NP} transforms into $-\phi^{SP}$, but also γ^6 goes into $-\gamma^6$, accordingly

$$\begin{aligned} \begin{matrix} 56 \\ (+) \end{matrix} & \text{ goes into } \begin{matrix} 56 \\ (-) \end{matrix} \\ \begin{matrix} 56 \\ [-] \end{matrix} & \text{ goes into } \begin{matrix} 56 \\ [+], \end{matrix} \end{aligned} \quad (43)$$

therefore $S^{56} \begin{matrix} 56 \\ (-) \end{matrix} = -\frac{1}{2} \begin{matrix} 56 \\ (-) \end{matrix}$ and $S^{56} \begin{matrix} 56 \\ [+], \end{matrix} = \frac{1}{2} \begin{matrix} 56 \\ [+]. \end{matrix}$ Taking into account Eqs. (43, 39, 32) we obtain

$$\begin{aligned} \psi_{n+\frac{1}{2}}^{SP(6)}(x^{NP\tau}) &= S \psi_{n+\frac{1}{2}}^{NP(6)}(x^{NP\tau}(x^{SP\tau})) \\ &= (i\alpha_n (\frac{(2\rho_0)^2}{\rho^{SP}}) e^{-i\phi^{SP}} \begin{matrix} 56 \\ (-) \end{matrix} \psi_{(+)}^{(4)} + \beta_n (\frac{(2\rho_0)^2}{\rho^{SP}}) \begin{matrix} 56 \\ [+], \end{matrix} \psi_{(-)}^{(4)}) e^{-in\phi^{SP}} \\ &= (i\alpha_{-(n+1)}^{SP} \begin{matrix} 56 \\ (-) \end{matrix} \psi_{(+)}^{(4)} + \beta_{-n}^{SP} e^{i\phi^{SP}} \begin{matrix} 56 \\ [+], \end{matrix} \psi_{(-)}^{(4)}) e^{-i(n+1)\phi^{SP}}. \end{aligned} \quad (44)$$

When evaluating $M^{SP56} = (S^{56} + i \frac{\partial}{\partial \phi^{SP}})$ on $S \psi_{n+\frac{1}{2}}^{NP(6)}(x^{NP\tau}(x^{SP\tau}))$ it follows

$$(S^{56} + i \frac{\partial}{\partial \phi^{SP}}) S \psi_{n+\frac{1}{2}}^{NP(6)}(x^{NP\tau}(x^{SP\tau})) = (n + \frac{1}{2}) S \psi_{n+\frac{1}{2}}^{NP(6)}. \quad (45)$$

Accordingly the massless state $\psi_{\frac{1}{2}}^{NP(6)m=0} = \mathcal{N}_0^{NP} f^{NP(-F+\frac{1}{2})} \begin{matrix} 56 \\ (+) \end{matrix} \psi_{(+)}^{(4)}$ from Eq. (21) looks, when transforming the coordinate system from the northern to the southern pole, as

$$\psi_{\frac{1}{2}}^{SP(6)m=0} = \mathcal{N}_0^{SP} (f^{SP} (\frac{2\rho_0}{\rho^{SP}})^2)^{(-F+\frac{1}{2})} \begin{matrix} 56 \\ (-) \end{matrix} \psi_{(+)}^{(4)} e^{-i\phi^{SP}}. \quad (46)$$

Taking into account that $x^{SP(5)} + i2S^{56}x^{SP(6)} = \rho^{SP} e^{-i2S^{56}\phi^{SP}}$ and $-\frac{\partial}{\partial x^{SP(5)}} + i2S^{56} - \frac{\partial}{\partial x^{SP(6)}} = e^{-i2S^{56}\phi^{SP}} (\frac{\partial}{\partial \rho^{SP}} - i2S^{56} \frac{1}{\rho^{SP}} \frac{\partial}{\partial \phi^{SP}})$ we can write the equations of motion as

$$\begin{aligned} if e^{-i\phi^{SP}2S^{56}} \{ (\frac{\partial}{\partial \rho^{SP}} - \frac{i2S^{56}}{\rho^{SP}} \frac{\partial}{\partial \phi^{SP}}) + S^{56} \frac{1}{\rho^{SP}} (\frac{4F}{f^{SP}} - 2 \cdot 2S^{56}) \\ + \frac{1}{\rho^{SP}} (1 - \frac{f^{SP} - 1}{f^{SP}}) \} \psi^{(6)} + \gamma^0 \gamma^5 m \psi^{(6)} = 0. \end{aligned} \quad (47)$$

For $\psi_{n+\frac{1}{2}}^{SP(6)} = (\mathcal{A}_{-(n+1)} e^{-i\phi^{SP}} \begin{matrix} 56 \\ (-) \end{matrix} \psi_{(+)}^{(4)} + \mathcal{B}_{-n} \begin{matrix} 56 \\ [+], \end{matrix} \psi_{(-)}^{(4)}) e^{-in\phi^{SP}}$ we find the equations for $\mathcal{A}_{-(n+1)}$ and $\mathcal{B}_{-n}$

$$\begin{aligned} -if \{ (\frac{\partial}{\partial \rho^{SP}} + \frac{-n}{\rho^{SP}}) + \frac{1}{\rho^{SP}} (\frac{2F+1}{f^{SP}} - 1) \} \mathcal{B}_{-n} + m \mathcal{A}_{-(n+1)} &= 0, \\ -if \{ (\frac{\partial}{\partial \rho^{SP}} + \frac{n+1}{\rho^{SP}}) + \frac{1}{\rho^{SP}} (\frac{-2F+1}{f^{SP}} - 1) \} \mathcal{A}_{-(n+1)} + m \mathcal{B}_{-n} &= 0. \end{aligned} \quad (48)$$

When using $f^{SP} \frac{\partial}{\partial \rho^{SP}} = \frac{1}{\rho_0} \frac{\partial}{\partial \vartheta^{SP}}$ and $\frac{f^{SP}}{\rho^{SP}} = \frac{1}{\rho_0} \frac{1}{\sin \vartheta^{SP}}$ Eq.(48) transforms into

$$\begin{aligned} \left(\frac{\partial}{\partial \vartheta^{SP}} + \frac{-n-1 + (F + \frac{1}{2})(1 + \cos \vartheta^{SP})}{\sin \vartheta^{SP}} \right) \mathcal{B}_{-n} + i\rho_0 m \mathcal{A}_{-(n+1)} &= 0. \\ \left(\frac{\partial}{\partial \vartheta^{SP}} + \frac{n + (-F + \frac{1}{2})(1 + \cos \vartheta^{SP})}{\sin \vartheta^{SP}} \right) \mathcal{A}_{-(n+1)} + i\rho_0 m \mathcal{B}_{-n} &= 0. \end{aligned} \quad (49)$$

Again we find for $2F = 1$

$$\begin{aligned} \left\{ \frac{1}{\sin \vartheta} \frac{\partial}{\partial \vartheta} (\sin \vartheta \frac{\partial}{\partial \vartheta}) + [(\rho_0 m)^2 - \frac{n^2}{\sin^2 \vartheta}] \right\} \mathcal{A}_{-(n+1)} &= 0, \\ \mathcal{B}_{-n} = i \frac{1}{(\rho_0 m)^2} \left(\frac{\partial}{\partial \vartheta^{SP}} + \frac{n}{\sin \vartheta^{SP}} \right) \mathcal{A}_{-(n+1)}. \end{aligned} \quad (50)$$

Let us conclude this section by recognizing that we have at the south pole allowed a certain special singularity which is of the following type: Around a point in the 2-dimensional space - the singular point - we let the phase of the wave function rotate so that it turns around 2π as one goes around 2π in the direction to the singular point *i.e.* as ϕ goes around. This would for a properly smooth function only be allowed provided that the magnitude of the wave function decreases linearly with the distance to the singular point. Of course, from the point of view of the structure of the singularity we can make a gauge transformation and replace the just mentioned phase rotation of the wave function by a singular (essentially constant) value of the spin connection on the circles around the singular point.

IV. SPINORS AND THE GAUGE FIELDS IN $d = (1 + 3)$

To study how do spinors couple to the Kaluza-Klein gauge fields in the case of $M^{(1+5)}$, “broken” to $M^{(1+3)} \times S^2$ with the radius of S^2 equal to ρ_0 and with the spin connection field $\omega_{st\sigma} = i4F\varepsilon_{st} \frac{x_\sigma}{\rho} \frac{f-1}{\rho f}$ we first look for (background) gauge gravitational fields, which preserve the rotational symmetry around the axis through the northern and southern pole. Requiring that the symmetry determined by the Killing vectors of Eq. (A.1) (following ref. [24]) with $f^\sigma_s = f\delta^\sigma_s, f^\mu_s = 0, e^s_\sigma = f^{-1}\delta^\sigma_s, e^m_\sigma = 0$, is preserved, we find for the background vielbein field

$$e^a_\alpha = \begin{pmatrix} \delta^m_\mu & e^m_\sigma = 0 \\ e^s_\mu & e^s_\sigma \end{pmatrix}, f^\alpha_a = \begin{pmatrix} \delta^\mu_m & f^\sigma_m \\ 0 = f^\mu_s & f^\sigma_s \end{pmatrix}, \quad (51)$$

with

$$\begin{aligned} f^\sigma_m &= K^{(56)\sigma} B^{(5)(6)}_\mu f^\mu_m = \varepsilon^\sigma_\tau x^\tau A_\mu \delta^\mu_m, \\ e^s_\mu &= -\varepsilon^\sigma_\tau x^\tau A_\mu e^s_\sigma, \end{aligned} \quad (52)$$

$s = 5, 6; \sigma = (5), (6)$. Requiring that correspondingly the only nonzero torsion fields are those from Eq. (A.2) we find for the spin connection fields

$$\omega_{st\mu} = \varepsilon_{st} A_\mu, \quad \omega_{sm\mu} = \frac{1}{2} f^{-1} \varepsilon_{s\sigma} x^\sigma \delta^\nu_m F_{\mu\nu}, \quad (53)$$

$F_{\mu\nu} = A_{[\nu, \mu]}$. The $U(1)$ gauge field A_μ depends only on x^μ . All the other components of the spin connection fields, except (by the Killing symmetry preserved) $\omega_{st\sigma}$ from Eq. (6), are zero, since for simplicity we allow no gravity in $(1+3)$ dimensional space. The corresponding nonzero torsion fields $\mathcal{T}^a_{bc}$ are presented in Eq. (A.2) and in the expressions following this equation, all the other components are zero.

To determine the current, which couples the spinor to the Kaluza-Klein gauge fields A_μ , we analyse (as in the refs. [24]) the spinor action (Eq. (6))

$$\begin{aligned} \mathcal{S} = & \int d^d x \bar{\psi}^{(6)} E \gamma^a p_{0a} \psi^{(6)} = \\ & \int d^d x \bar{\psi}^{(6)} \gamma^s p_s \psi^{(6)} + \\ & \int d^d x \bar{\psi}^{(6)} \gamma^m \delta^\mu_m p_\mu \psi^{(6)} + \\ & \int d^d x \bar{\psi}^{(6)} \gamma^m \delta^\mu_m A_\mu (\varepsilon^\sigma_\tau x^\tau p_\sigma + S^{56}) \psi^{(6)} + \\ & \text{terms } \propto x^\sigma \text{ or } \propto x^5 x^6. \end{aligned} \quad (54)$$

Here $\psi^{(6)}$ is a spinor state in $d = (1+5)$ after the break of M^{1+5} into $M^{1+3} \times S^2$. E is for f^α_a from Eq. (51) equal to f^{-2} . The term in the second row in Eq. (54) is the mass term (equal to zero for the massless spinor), the term in the third row is the kinetic term, together with the term in the fourth row defines the covariant derivative $p_{0\mu}$ in $d = (1+3)$. The terms in the last row contribute nothing when the integration over the disk (curved into a sphere S^2) is performed, since they all are proportional to x^σ or to $\varepsilon_{\sigma\tau} x^\sigma x^\tau$ ($-\gamma^m \frac{1}{2} S^{sm} \omega_{smn} = -\gamma^m \frac{1}{2} f^{-1} F_{mn} \varepsilon_{s\sigma} x^\sigma$ and $-\gamma^m f^\sigma_m \frac{1}{2} S^{st} \omega_{st\sigma} = \gamma^m A_m x^5 x^6 S^{st} \varepsilon_{st} \frac{4iF(f-1)}{f\rho^2}$).

We end up with the current in $(1+3)$

$$j^\mu = \int E d^2 x \bar{\psi}^{(6)} \gamma^m \delta^\mu_m M^{56} \psi^{(6)}. \quad (55)$$

The charge in $d = (1+3)$ is proportional to the total angular momentum $M^{56} = L^{56} + S^{56}$ around the axis from the southern to the northern pole of S^2 , but since for the choice of $2F = 1$ (and for any $0 < 2F \leq 1$) in Eq. (19) only a left handed massless spinor exists, with the angular momentum zero, the charge of a massless spinor in $d = (1+3)$ is equal to $1/2$.

The Riemann scalar is for the vielbein of Eq. (51) equal to $\mathcal{R} = -\frac{1}{2}\rho^2 f^{-2} F^{mn} F_{mn}$.

If we integrate the Riemann scalar over the fifth and the sixth dimension, we get $-\frac{8\pi}{3}(\rho_0)^4 F^{mn} F_{mn}$.

V. CONCLUSIONS

We prove in this paper that one can escape from the "no-go theorem" of Witten [15], that is one can guarantee the masslessness of spinors and their chiral coupling to the Kaluza-Klein[like] gauge fields when breaking the symmetry from d -dimensional one to $M^{(1+3)} \times M^{d-4}$ space, in cases which we call the "effective two dimensionality" even without boundaries, as we proposed in the references [24]. Namely, we can guarantee the above mentioned properties of spinors, when the break $M^{(1+3)} \times M^{d-4}$, $d - 4 > 2$ occurs in a way that vielbeins and spin connections are completely flat in all but two dimensions, while the two dimensional space, although of finite volume, is noncompact with a particular spin connection contributing to the properties of spinors. In our particular case it is the zweibein (the zweibein of the S^2 sphere with a hole at the southern pole) on an infinite disc, which guarantees that the noncompact space has the finite volume and enables, together with the spin connection field on this disc ($\omega_{st\sigma} = i F \varepsilon_{st} \frac{x_\sigma}{f \rho_0^2}$, the $\omega_{st\sigma}$ field breaks the parity symmetry and the sign of F makes a choice of the handedness of the massless state) that only one normalizable spinor state (of particular handedness) is massless, carrying the Kaluza-Klein charge of $\frac{1}{2}$ and coupling chirally to the corresponding Kaluza-Klein gauge field. Let us add that requiring normalizability of states in extra dimensions guarantees that states are normalizable in the whole $d = (1 + (d - 1))$ space.

Since the spin connection strength is determined only within an interval ($0 < 2F \leq 1$), what we proposed is not a fine tuning. Taking (in the absence of fermions) the action for the gravitational gauge fields with the linear curvature for $d = 2$ (when any zweibein and any spin connection fulfills the corresponding equations of motion), we are allowed to make any choice of a zweibein and spin connection. (This choice leads to nonzero torsion.)

There is the discrete spectrum of normalizable eigenstates of the Hermitean Hamiltonian on the infinite disc for the chosen zweibein and spin connection of any strength F in the interval ($0 < 2F \leq 1$), as we proved in section II.

The normalizable eigenstates, which are chosen to be at the same time the eigenstates

of the total angular momentum on the disc $M^{56} = x^5 p^6 - x^6 p^5 + S^{56}$, with the eigenvalues $(n + 1/2)$, carry the Kaluza-Klein charge $(n + 1/2)$. The only massless state carries the charge $(\frac{1}{2})$. For the choice of $2F = 1$ the normalizable massless state is independent of the coordinates on the disc. The normalizable massive states have the masses equal to $k(k + 1)/\rho_0$, $k = 1, 2, 3, \dots$, with $-k \leq n \leq k$. The spectrum is obviously discrete and stays discrete for all F in the interval $0 < 2F \leq 1$ and for any finite ρ_0 . The current is for all the solutions and also for all F equal to zero. As long as the Hamiltonian is Hermitean on a disc, fermions can not leave the disc, unless an additional interaction (or a dynamical restoration of the symmetry $M^{(1+5)}$, that is the *phase transition*) would force them to go out of the disc, which is not the case for our toy model.

Understanding the infinite disc as a S^2 sphere with the southern pole missing, a singularity of the type should be recognized: Around a point in the 2-dimensional space of S^2 - the singular point - we let the phase of the wave function rotate so that it turns around 2π as one goes around 2π in the direction to the singular point. But from the point of view of the structure of the singularity we can make a gauge transformation and replace the just mentioned phase rotation of the wave function by a singular (essentially constant) value of the spin connection on the circles around the singular point.

The possibility that after the break a two dimensional manifold (with the zweibein of S^2 , with one point missing and with a particular spin connection field) exists allowing only one normalizable massless state which is correspondingly mass protested and which couples to the Kaluza-Klein charge, opens, to our understanding, a new hope for the Kaluza-Klein[like] theories of the elegant version, with only the gravity, and will help to revive them.

Appendix: The Killing vectors and the torsion terms for our model

The infinitesimal coordinate transformations manifesting the symmetry of M^{1+3} and the S^2 are: $x'^\mu = x^\mu$, $x'^\sigma = x^\sigma + \phi_A K^{A\sigma}$, with ϕ_A the parameter of rotations around the axis which goes through both poles and with the infinitesimal generators of rotations around this axis $M^{(5)(6)} (= x^{(5)} p^{(6)} - x^{(6)} p^{(5)} + S^{(5)(6)})$

$$K^{A\sigma} = K^{(56)\sigma} = -iM^{(5)(6)} x^\sigma = \varepsilon^\sigma{}_\tau x^\tau, \quad (\text{A.1})$$

with $\varepsilon^\sigma{}_\tau = -1 = -\varepsilon_\tau{}^\sigma$, $\varepsilon^{(5)(6)} = 1$. The operators $K_\sigma^A = f^{-2}\varepsilon_{\sigma\tau}x^\tau$ fulfil the Killing relation $K_{\sigma,\tau}^A + \Gamma^{\sigma'}{}_{\sigma\tau}K_{\sigma'}^A + K_{\tau,\sigma}^A + \Gamma^{\sigma'}{}_{\tau\sigma}K_{\sigma'}^A = 0$, (with $\Gamma^{\sigma'}{}_{\sigma\tau} = -\frac{1}{2}g^{\rho\sigma'}(g_{\tau\rho,\sigma} + g_{\sigma\rho,\tau} - g_{\sigma\tau,\rho})$).

From $\gamma^ap_{0a}\gamma^bp_{0b} = p_{0a}p_0^a - iS^{ab}S^{cd}\mathcal{R}_{abcd} + S^{ab}\mathcal{T}^\beta{}_{ab}p_{0\beta}$ we find for the torsion

$$\mathcal{T}^\beta{}_{ab} = f^\alpha{}_{[a}(f^\beta{}_{b]})_{,\alpha} + \omega_{[a}{}^c{}_{b]}f_c{}^\beta. \quad (\text{A.2})$$

From Eq. (A.2) we read that to the torsion on S^2 both, the zweibein f_τ^σ and the spin connection $\omega_{st\sigma}$, contribute. While we have on S^2 for $\mathcal{R}_{\sigma\tau} = f^{-2}\eta_{\sigma\tau}\frac{1}{\rho_0^2}$ and correspondingly for the curvature $\mathcal{R} = \frac{-2}{(\rho_0)^2}$, we find for the torsion $\mathcal{T}^s{}_{ts'} = \mathcal{T}^s{}_{t\sigma}f_{s'}^\sigma$ with $\mathcal{T}^5{}_{ss} = 0 = \mathcal{T}^6{}_{ss}$, $s = 5, 6$, $\mathcal{T}^5{}_{65} = -\mathcal{T}^5{}_{56} = -(f_{,6} + \frac{4iF(f-1)}{\rho^2}x_5)$, $\mathcal{T}^6{}_{56} = -\mathcal{T}^6{}_{65} = -f_{,5} + \frac{4iF(f-1)}{\rho^2}x_6$. The torsion $\mathcal{T}^2 = \mathcal{T}^s{}_{ts'}\mathcal{T}^{ts'}$ is for our particular choice of the zweibein and spin connection fields from Eqs. (3,3) correspondingly equal to $-\frac{2\rho^2}{(\rho_0)^4}(1 - (2F)^2)$.

-
- [1] T. Kaluza, *Sitzungsber.Preuss.Akad.Wiss.Berlin, Math.Phys.* **K1** (1921) 966, O. Klein, *Z.Phys.* **37** (1926) 895.
 - [2] H. Georgi, S. L. Glashow, *Phys. Rev. Lett.* **32** (1974) 438.
 - [3] Y. M. Cho, *J. Math. Phys.* **16** (1975) 2029, Y. M. Cho, P. G. O.Freund, *Phys. Rev.* **D 12** (1975) 1711.
 - [4] A. Zee, *Proceedings of the first Kyoto summer institute on grand unified theories and related topics*, Kyoto, Japan, June-July 1981, Ed. by M. Konuma, T. Kaskawa, World Scientific Singapore.
 - [5] A. Salam, J. Strathdee, *Ann. Phys. (N.Y.)* **141** (1982) 316.
 - [6] S. Randjbar-Daemi, A. Salam, J. Strathdee, *Nucl. Phys.* **B 242** (1984) 447.
 - [7] W. Mecklenburg, *Fortschr. Phys.* **32** (1984) 207.
 - [8] M. Duff, B. Nilsson, C. Pope, *Phys. Rep.* **C 130** (1984)1, M. Duff, B. Nilsson, C. Pope, N. Warner, *Phys. Lett.* **B 149** (1984) 60.
 - [9] Z. Horvath, L. Palla, E. Crammer, J. Scherk, *Nucl. Phys.* **B 127** (1977) 57.
 - [10] G. Chapline, R. Slansky, *Nucl. Phys.* **B 209** (1982) 461.
 - [11] T. Appelquist, H. C. Cheng, B. A. Dobrescu, *Phys. Rev.* **D 64** (2001) 035002, I. Antoniadis, *Phys. Lett.* **B 246** (1990) 377.

- [12] M. Saposhnikov, P. Tinyakov, *Phys. Lett.* **B 515** (2001) 442, arxiv:hep-th/0102161v2, S. Randjbar-Daemi, M. Saposhnikov, *Nucl. Phys.* **B 645** (2002) 188.
- [13] C. Wetterich, *The second Jerusalem Winter school on theoretical physics*, 1985, CERN-TH4190/85 and references therein, *Nucl. Phys.* **B 253** (1985) 261, *Nucl. Phys.* **B 234** (1984) 413.
- [14] The authors of the works presented in *An introduction to Kaluza-Klein theories*, Ed. by H. C. Lee, World Scientific, Singapore 1983.
- [15] E. Witten, *Nucl. Phys.* **B 186** (1981) 412; Princeton Technical Rep. PRINT -83-1056, October 1983.
- [16] N. Arkani-Hamed, S. Dimopoulos, G. Dvali, *Phys. Lett.* **B429** (1998) 263, arXiv:hep-ph/9803315v1; *Phys. Rev.* **D 59** (1999) (1998) 086004, arXiv:hep-ph/9807344.
- [17] V.A. Rubakov, *Phys. Usp.* **44** (2001) 871; *Usp. Fiz. Nauk* **171** (2001) 913, arXiv:hep-ph/0104152v2.
- [18] L. Randall and R. Sundrum, *Phys. Rev. Lett.* **83** (1999) 3370, [hep-ph/9905221]; *Phys. Rev. Lett.* **83** (1999) 4690, [hep-th/9906064].
- [19] M. Blagojević, *Gravitation and Gauge Symmetries*, IOP Publishing, Bristol, 2001, ISBN-10: 0750307676, ISBN-13: 978-0750307673.
- [20] S. Deser, gr-qc/9512022.
- [21] N. S. Mankoč Borštnik, *Phys. Lett.* **B 292** (1992) 25, *J. Math. Phys.* **34** (1993) 3731 Int. J. Theor. Phys. **40** 315 (2001), *Modern Phys. Lett.* **A 10** (1995) 587, A. Borštnik, N. S. Mankoč Borštnik, hep-ph/0401043, hep-ph/0401055, hep-ph/0301029, *Phys. Rev.* **D 74** (2006)073013, hep-ph/0512062, G.regar, M. Breskvar, D. Lukman, N. S. Mankoč Borštnik, hep-ph/0711.4681, *New J. of Phys.* **10** (2008) 093002, hep-ph/0606159, hep-ph/07082846, hep-ph/0612250, p.25-50.
- [22] N. S. Mankoč Borštnik, <http://arxiv.org/abs/1011.5765>, <http://arxiv.org/abs/1011.5764>, and talks and discussions in <http://arxiv.org/abs/1012.0224>.
- [23] N. Nicolai, C. Wetterich, *Phys. Lett.* **B 150** (1985)347.
- [24] N. S. Mankoč Borštnik, H. B. Nielsen, *Phys. Lett.* **B 633** (2006) 771-775, hep-th/0311037, hep-th/0509101, *Phys. Lett.* **B 644** (2007)198-202, hep-th/0608006, *Phys. Lett.* **B 10.1016** (2008), N. S. Mankoč Borštnik, H. B. Nielsen, D. Lukman, hep-ph/0412208.
- [25] N. S. Mankoč Borštnik, H. B. Nielsen, *J. of Math. Phys.* **43** (2002) 5782, hep-th/0111257, *J.*

of *Math. Phys.* **44** (2003) 4817, hep-th/0303224.

- [26] Z. X. Wang, D. R. Guo, *Special functions*, World Scientific, 1989, sect. 5.18., pages 255-258.
- [27] The author of the ref. [13] proposes, for example, the "squashed" S^2 sphere, recognizing that with the zweibein of S^2 (he calls in this case S^2 a compact space) there are no massless spinor states, while with at least a little "stronger" zweibein than with that of S^2 (like with $f = (1 + (\frac{\rho}{2\rho_0})^{2+k})$, with $0 < k \leq 2$, $k = 0$ reproduces S^2) there are two massless states. Although the author wrote differently, these two massless states belong to the left and the right handed state with respect to $d = (1 + 3)$, and therefore not mass protected, and would correspondingly lead to massive fermion states.
- [28] In the ref. [13], mentioned and discussed in the previous footnote, this idea of a finite volume of a noncompact space, as well as the normalizability of states is already stressed.
- [29] The *approach unifying spin and charges and predicting families* [21] proposes in $d = (1 + (d-1))$ a simple starting action for spinors with the two kinds of the spin generators (γ matrices): the Dirac one, which takes care of the spin and the charges, and the second one, anticommuting with the Dirac one, which generates families. For the explanation of the appearance of the two kinds of the spin generators we invite the reader to look at the refs. [21, 25] and the references therein. A spinor couples in $d = 1 + 13$ to the vielbeins and (through two kinds of the spin generators to) the spin connection fields. Appropriate breaks of the starting symmetry lead to the left handed quarks and leptons in $d = (1 + 3)$, which carry the weak charge while the right handed ones are weak chargeless. The *approach* is offering the answers to the questions about the origin of families of quarks and leptons, about the explicit values of their masses and mixing matrices (predicting the fourth family to be possibly seen at the LHC or at somewhat higher energies) as well as about the masses of the scalar and the weak gauge fields, about the dark matter candidates, and about breaking the discrete symmetries. There are many possibilities in the *approach* for breaking the starting symmetries to those of the *standard model*. These problems were studied in some crude approximations in refs. [21] and are under consideration [22].
- [30] We have proven in ref. [24] that only in even dimensional spaces of $d = 2$ modulo 4 dimensions (*i.e.* in $d = 2(2n + 1)$, $n = 0, 1, 2, \dots$) spinors (they are allowed to be in families) of one handedness and with no conserved charges gain no Majorana mass.
- [31] f^α_a are inverted vielbeins to e^α_a with the properties $e^\alpha_a f^\alpha_b = \delta^a_b$, $e^\alpha_a f^\beta_a = \delta^\beta_\alpha$. Latin

indices $a, b, \dots, m, n, \dots, s, t, \dots$ denote a tangent space (a flat index), while Greek indices $\alpha, \beta, \dots, \mu, \nu, \dots, \sigma, \tau, \dots$ denote an Einstein index (a curved index). Letters from the beginning of both the alphabets indicate a general index ($a, b, c, \dots$ and $\alpha, \beta, \gamma, \dots$), from the middle of both the alphabets the observed dimensions 0, 1, 2, 3 ($m, n, \dots$ and $\mu, \nu, \dots$), indices from the bottom of the alphabets indicate the compactified dimensions ($s, t, \dots$ and $\sigma, \tau, \dots$). We assume the signature $\eta^{ab} = \text{diag}\{1, -1, -1, \dots, -1\}$.

- [32] One finds that $\omega_{cda} = \Re \omega_{cda}$, if c, d, a all different while $\omega_{cda} = i \Im \omega_{cda}$, otherwise.
- [33] It is expected that the zweibein curving the infinite disc into an (almost S^2) and the spin connection, which breaks the parity symmetry and takes a part in determining equations of motion, appear dynamically, causing the "phase transition". Accordingly could dynamical fields by causing the phase transition restore the symmetry of M^{1+5}
- [34] Rewriting the mass operator $\hat{m} = \gamma^0 \gamma^s f^\sigma_s (p_\sigma - S^{56} \omega_{56\sigma} + \frac{1}{2Ef} \{p_\sigma, Ef\}_-)$ as a function of ϑ and ϕ : $\rho_0 \hat{m} = i \gamma^0 \{ \overset{56}{(+)} e^{-i\phi} (\frac{\partial}{\partial \vartheta} - \frac{i}{\sin \vartheta} \frac{\partial}{\partial \phi} - \frac{1-\cos \vartheta}{\sin \vartheta}) + \overset{56}{(-)} e^{i\phi} (\frac{\partial}{\partial \vartheta} + \frac{i}{\sin \vartheta} \frac{\partial}{\partial \phi}) \}$, one can easily show that when applying $\rho_0 \hat{m}$ and M^{56} on $\psi_{n+1/2}^{(6)\hat{m}^2=l(l+1)}$, for $l = 1, 2, \dots$, one obtains from Eq. (24) $\rho_0 \hat{m} \psi_{n+1/2}^{(6)\hat{m}^2=l(l+1)} = l(l+1) \psi_{n+1/2}^{(6)\hat{m}^2=l(l+1)}$, $M^{56} \psi_{n+1/2}^{(6)\hat{m}^2=l(l+1)} = (n+1/2) \psi_{n+1/2}^{(6)\hat{m}^2=l(l+1)}$, $l = 1, 2, \dots$. A wave packet, which is the eigen function of M^{56} with the eigenvalue 1/2, for example, can be written as $\psi_{1/2}^{(6)} = \sum_{k=0, \infty} C_{1/2}^k \mathcal{N}_{1/2} \{ \overset{56}{(+)} \psi_{(+)}^{(4)} + (1 - \delta_0^k) \frac{i}{\sqrt{k(k+1)}} \overset{56}{[-]} \psi_{(-)}^{(4)} e^{i\phi} \frac{\partial}{\partial \vartheta} \} Y_0^k$. The expectation value of the mass operator $\hat{m}$ on such a wave packet is $\sum_{k=0, \infty} C_{1/2}^{k*} C_{1/2}^k \sqrt{k(k+1)} / \rho_0$.